

# Applications of semidefinite programming to combinatorial problems

---

Norihide Tokushige (University of the Ryukyus)

Spectral graph theory and related topics

January 26–28th, 2025 @Waseda University

# Outline

- References and Software for SDP
- Applications of SDP for intersecting families
  - A toy problem (independence number of the Petersen graph)
  - Wilson's proof of Erdős–Ko–Rado Theorem
  - Recent results on cross intersecting families

# References

## Applications of SDP/LP for some concrete problems

- Wagner, Adam Zsolt. Refuting conjectures in extremal combinatorics via linear programming. JCTA (2020)
- Schrijver, Alexander. New code upper bounds from the Terwilliger algebra and semidefinite programming. IEEE (2005)
- 田中太初. Terwilliger 代数に基づく符号の半正定値計画限界. 代数学シンポ (2006)
- Bansal, Nikhil. Constructive algorithms for discrepancy minimization. FOCS (2010)

## Semidefinite programming

- M. J. Todd.  
Semidefinite optimization.  
Acta Numer. 10 (2001) 515–560.
- B. Gärtner, J. Matoušek.  
Approximation algorithms and semidefinite programming.  
Springer, 2012. xii+251 pp.

## Intersecting families / Association Schemes

- Godsil–Meagher. Erdős–Ko–Rado theorems: algebraic approaches. Cambridge Stud. Adv. Math., 2016.

# Software

## SDP solver

- SDPA on NEOS server



- SDPA の使い方 (How to start using SDPA)

## An example

- Consider the following primary program:

minimize  $\alpha$

$$\text{subject to } S := - \begin{bmatrix} \frac{9}{16} & \frac{3}{16} \\ \frac{3}{16} & \frac{1}{16} \end{bmatrix} + \alpha \begin{bmatrix} \frac{3}{4} & 0 \\ 0 & \frac{1}{4} \end{bmatrix} + \begin{bmatrix} x_1 & x_2 \\ x_2 & 0 \end{bmatrix} - \begin{bmatrix} z_1 & z_2 \\ z_2 & z_3 \end{bmatrix},$$
$$S \succeq 0, \quad z_1, z_2, z_3 \geq 0.$$

- true value for  $\min \alpha$  is  $\frac{1}{4}$ .

```
6
2
(2,-3)
{1,0,0,0,0,0}
0 1 1 1 0.5625
0 1 1 2 0.1875
0 1 2 2 0.0625
1 1 1 1 0.75
1 1 2 2 0.25
2 1 1 1 1
3 1 1 2 1
4 1 1 1 -1
4 2 1 1 1
5 1 1 2 -1
5 2 2 2 1
6 1 2 2 -1
6 2 3 3 1
```

```
phase.value = dFEAS
  Iteration = 35
            mu = 6.4453585130804993e-18
relative gap = 1.3840612359273038e-17
          gap = 3.2226792565402498e-17
        digits = 1.6256784703351158e+01
objValPrimal  = 2.5000000000000000e-01
objValDual    = 2.5000000000000000e-01
p.feas.error  = 4.3368086899420177e-19
d.feas.error  = 3.3787568866817071e-31
relative eps  = 4.9303806576313200e-32
total time    = 0.000
```

# Applications of SDP to intersecting families

- $[n] = \{1, 2, \dots, n\}$ .
- $\binom{[n]}{k} = \{F \subset [n] : |F| = k\}$ . (the set of  $k$ -element subsets)
- $\mathcal{F} \subset \binom{[n]}{k}$  is intersecting if  $F \cap F' \neq \emptyset$  for all  $F, F' \in \mathcal{F}$ .
- $\mathcal{F} \subset \binom{[n]}{k}$  is  $t$ -intersecting if  $|F \cap F'| \geq t$  for all  $F, F' \in \mathcal{F}$ .

**What is the maximum size of a  $t$ -intersecting family  $\mathcal{F} \subset \binom{[n]}{k}$  ?**

- The answer is known. The complete intersection theorem by Ahlswede and Khachatrian (1995).
- Wilson (1983) proved one of the main cases by solving an SDP.



## Cross $t$ -intersecting families

- Let  $\mathcal{F}, \mathcal{G} \subset \binom{[n]}{k}$ .
- $\mathcal{F}$  and  $\mathcal{G}$  are cross  $t$ -intersecting if  $|F \cap G| \geq t$  for  $\forall F \in \mathcal{F}, G \in \mathcal{G}$ .
- What is  $\max |\mathcal{F}| |\mathcal{G}|$  ?

Theorem (Zhang–Wu arXiv:2410.22792)

Let  $n \geq (t+1)(k-t+1)$ ,  $t \geq 3$ .

Under above conditions we have  $|\mathcal{F}| |\mathcal{G}| \leq \binom{n-t}{k-t}^2$ .

We settled the case  $t = 2$  by solving an SDP problem.

Theorem. The above result is also true for  $t = 2$ .


The case  $t = 1$  was settled by Pyber (1986), T (2013).

# A toy problem

A family  $\mathcal{F}$  is intersecting if  $F \cap F' \neq \emptyset$  for all  $F, F' \in \mathcal{F}$ .


What is the maximum size of an intersecting family  $\mathcal{F} \subset \binom{[5]}{2}$ ?

- $\binom{[5]}{2} = \{12, 13, 14, 15, 23, 24, 25, 34, 35, 45\}$ . (I write 12 for  $\{1, 2\}$ .)
- examples of intersecting families in  $\binom{[5]}{2}$ :




What is the maximum size of an intersecting family  $\mathcal{F} \subset \binom{[5]}{2}$  ?

- Kneser graph  $G = G(5, 2)$
- $V(G) = \binom{[5]}{2}$
- $x \sim y \iff x \cap y \neq \emptyset$ .



What is the maximum size of an intersecting family  $\mathcal{F} \subset \binom{[5]}{2}$  ?

- $U \subset V(G)$  is independent if no edges inside  $U$ .
- $\text{indep}(G) := \max\{|U| : U \text{ is independent}\}.$
- $\max |\mathcal{F}| = \text{indep}(G) \geq 4.$



## What is $\text{indep}(G)$ ?

- adjacency matrix  $A$  of  $G$ :  $(A)_{x,y} = \begin{cases} 1 & \text{if } x \sim y, \\ 0 & \text{if } x \not\sim y. \end{cases}$
- For  $G = G(5, 2)$ ,

$$A = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

- eigenvalues are  $3, 1, -2$ .

## positive semidefinite matrix

- Let  $M$  be an  $n \times n$  real symmetric matrix.
- $M$  is positive semidefinite if  $x^T M x \geq 0$  for all  $x \in \mathbb{R}^n$ .
- We write  $M \succeq 0$  if  $M$  is positive semidefinite.
- $M \succeq 0$  iff all eigenvalues are non-negative.
- For two matrices  $A, B$ , let  $A \bullet B = \sum_{x,y} (A)_{x,y} (B)_{x,y}$ .  
ex.  $\begin{bmatrix} a & b \\ b & c \end{bmatrix} \bullet \begin{bmatrix} x & y \\ y & z \end{bmatrix} = ax + 2by + cz$ .
- $x^T M x = M \bullet (xx^T)$ .
- If  $M \succeq 0$  then  $M \bullet (xx^T) \geq 0$ .

## Bounding indep( $G$ )

- Let  $U \subset V(G)$  be an independent set in  $G$ .
- Let  $\mathbf{u} \in \{0, 1\}^{10}$  be the indicator (column) vector of  $U$ , that is,

$$(\mathbf{u})_x = \begin{cases} 1 & \text{if } x \in U, \\ 0 & \text{if not.} \end{cases}$$

- Let  $X = \mathbf{u}\mathbf{u}^\top$ . Then  $(X)_{x,y} = 0$  if  $x \sim y$ .

## Bounding $\text{indep}(G)$

- Let  $U \subset V(G)$  be an independent set in  $G$ .
- Let  $\mathbf{u}$  be the indicator of  $U$ , and let  $X = \mathbf{u}\mathbf{u}^T$ .
- Let  $A$  be the adjacency matrix of the Kneser graph  $G = G(5, 2)$ .
- $I \bullet X = |U|$ .
- $J \bullet X = |U|^2$ .
- $A \bullet X = 0$ , that is,  $(A)_{x,y} (X)_{x,y} = 0$  for all  $x, y$ .
- Let  $S := \alpha I - J + \beta A$ . Then  $S \bullet X = \alpha|U| - |U|^2$ .
- If  $S \succeq 0$  then  $S \bullet X \geq 0$ , so  $|U| \leq \alpha$ , i.e.,  $\text{indep}(G) \leq \alpha$ .



## What is $\text{indep}(G)$ ?

- Let  $S := \alpha I - J + \beta A$ .
- If  $S \succeq 0$ , then  $\text{indep}(G) \leq \alpha$ .
- For  $G = G(5, 2)$ , let  $S = 4I - J + 2A$ . Then  $S \succeq 0$ .
- That is,  $\text{indep}(G) \leq 4$ . (So,  $\text{indep}(G) = 4$ )

## An SDP problem for $\text{indep}(G(5, 2))$

minimize  $\alpha$

subject to  $S := \alpha I - J + \beta A \succeq 0$ . (variables are  $\alpha, \beta$ .)

- A feasible solution  $\alpha$  satisfies  $\text{indep}(G) \leq \alpha$ .

## Extending adjacency matrix

- To bound  $\text{indep}(G)$  we used  $A \bullet X = 0$ .
- For this, we didn't use  $(A)_{x,y} = 1$  if  $x \sim y$ .
- This means that if  $x \sim y$ , then  $(A)_{x,y}$  is not necessarily 1.
- Redefine an “adjacency matrix” by

$$(A)_{x,y} = \begin{cases} 0 & \text{if } x \not\sim y, \\ * & \text{if not,} \end{cases}$$

where  $*$  is any number (provided  $A$  is symmetric).

# Bounding the size of $t$ -intersecting families in $\binom{[n]}{k}$

- $\mathcal{F} \subset \binom{[n]}{k}$  is  $t$ -intersecting if  $|F \cap F'| \geq t$  for all  $F, F' \in \mathcal{F}$ .
- $\mathcal{F} = \{F \in \binom{[n]}{k} : [t] \subset F\}$  is  $t$ -intersecting, and  $|\mathcal{F}| = \binom{n-t}{k-t}$ .
- What is the maximum size of  $t$ -intersecting families  $\mathcal{F} \subset \binom{[n]}{k}$  ?
- Kneser graph  $G = G(n, k, t)$ :  $V(G) = \binom{[n]}{k}$ ,  $x \sim y \iff |x \cap y| < t$ .
- What is  $\text{indep}(G)$  ? By construction,  $\text{indep}(G) \geq \binom{n-t}{k-t}$ .
- (An SDP problem) **minimize  $\alpha$**   
**subject to  $S := \alpha I - J + A \succeq 0$ , where  $(A)_{x,y} = 0$  if  $x \not\sim y$ .**
- A feasible solution  $\alpha$  satisfies  $\text{indep}(G) \leq \alpha$ .
- Wilson found an  $S \succeq 0$  with  $\alpha = \binom{n-t}{k-t}$  if  $n$  is not too small.

- Let  $n \geq (t+1)(k-t+1)$ .
- Wilson found an  $A$  satisfying  $(A)_{x,y} = 0$  for  $|x \cap y| \geq t$  and

$$S = \binom{n-t}{k-t} I - J + A \succeq 0.$$

- ex. Wilson's matrix for  $G(8, 3, 2)$ :  $(A)_{x,y} = \begin{cases} \frac{1}{2} & \text{if } |x \cap y| = 0, \\ \frac{3}{2} & \text{if } |x \cap y| = 1, \\ 0 & \text{if } |x \cap y| \geq 2. \end{cases}$
- This implies that  $\text{indep}(G) \leq \binom{n-t}{k-t}$ , so  $\text{indep}(G) = \binom{n-t}{k-t}$ .

Theorem. Let  $n \geq (t+1)(k-t+1)$ .

If  $\mathcal{F} \subset \binom{[n]}{k}$  is  $t$ -intersecting, then  $|\mathcal{F}| \leq \binom{n-t}{k-t}$ .

# Cross 2-intersecting families

- $\mathcal{F}$  and  $\mathcal{G}$  are cross 2-intersecting if  $|F \cap G| \geq 2$  for  $\forall F \in \mathcal{F}, G \in \mathcal{G}$ .
- If  $\mathcal{F} = \mathcal{G} = \{F \in \binom{[n]}{k} : \{1, 2\} \subset F\}$ , then they are cross 2-intersecting, and  $|\mathcal{F}| = |\mathcal{G}| = \binom{n-2}{k-2}$ .

We got the following result by solving an SDP problem.

Theorem. Let  $\mathcal{F}, \mathcal{G} \subset \binom{[n]}{k}$  and  $n \geq 3(k-1)$ .

If  $\mathcal{F}$  and  $\mathcal{G}$  are cross 2-intersecting, then  $|\mathcal{F}||\mathcal{G}| \leq \binom{n-2}{k-2}^2$ .

## SDP for Cross 2-intersecting families


- (Suda–Tanaka 2014) minimize  $\alpha$ , subject to

$$S := \frac{1}{2} \begin{bmatrix} \alpha I & -J \\ -J & \alpha I \end{bmatrix} + A - Z \succeq 0, \quad Z \succeq 0, \quad (A)_{x,y} = 0 \text{ for } |x \cap y| \geq 2.$$

- (★) A feasible solution  $\alpha$  satisfies  $|\mathcal{F}||\mathcal{G}| \leq \alpha^2$ .
- If  $n \geq 3(k-1)$ , then we can find  $A$  and  $Z \succeq 0$  so that  $S \succeq 0$  with  $\alpha = \binom{n-2}{k-2}$ .
- What is this  $Z$  anyway?

(talk at RIMS, Kyoto University, 6th March)

# Bipartite Kneser graph



## Conjecture

Let  $k \geq l$  and  $n \geq 3(k - 1)$ . Suppose that  $\mathcal{F} \subset \binom{[n]}{k}$  and  $\mathcal{G} \subset \binom{[n]}{l}$  are cross 2-intersecting. Then  $|\mathcal{F}||\mathcal{G}| \leq \binom{n-2}{k-2} \binom{n-2}{l-2}$ .

## Problems

- families of subspaces in a vector space
- families of permutations
- other structures, e.g., partitions, perfect matching, etc.
- See also Godsil–Meagher (Chapter 16, Open problems).